

ORAL PRESENTATION

Open Access

Uncovering a novel function of BTLA on tumor-infiltrating CD8+ T cells

Cara Haymaker^{1*}, Richard Wu¹, Krit Ritthipichai¹, Chantale Bernatchez¹, Marie-Andree Forget¹, Jie Qing Chen¹, Hiu Liu², Ena Wang², Francesco Marincola^{2,3}, Patrick Hwu¹, Laszlo Radvanyi¹

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

Manipulation of T-cell co-inhibitory molecules, such as CTLA-4, PD-1, and BTLA has recently moved to the forefront of cancer immunotherapy. Although these molecules serve as inhibitors of T-cell activation, they are also biomarkers for activated T cells and may in fact have positive immune regulatory functions under certain circumstances. Recently, we demonstrated an unexpected positive association of CD8+ T cells expressing BTLA (B- and T- lymphocyte attenuator) with clinical response to adoptive T cell therapy in late-stage melanoma patients. We hypothesized that TIL may utilize the BTLA checkpoint differently and that key phenotypic and functional differences may exist between CD8+BTLA+ and CD8+BTLA- TIL subsets. In this study, we isolated and characterized BTLA+ and BTLA- CD8+ TIL from melanoma patients accrued in a Phase II clinical trial. We found that CD8+BTLA+ TIL had a superior proliferative response to IL-2 and a younger, more central memory T-cell behavior, such as secreting their own IL-2 after TCR stimulation. This younger and more robust memory phenotype was also associated with a longer persistence of T-cell clones in vivo in patients from the infused CD8+BTLA+ TIL subset. In contrast, CD8+BTLA- TIL were poorly proliferative, expressed killer-cell immunoglobulin-like receptors, and exhibited a gene expression signature of T cell deletion. As previously demonstrated, BTLA ligation with its cognate ligand herpes virus entry mediator (HVEM) resulted in decreased proliferation and inflammatory cytokine secretion. However, in a model of TIL activation induced cell death, we observed an enhanced survival of TIL co-cultured with HVEM+ target cells suggesting that BTLA ligation may also promote T-cell survival. Using a recombinant HVEM-Fc protein, we found that HVEM

ligation of BTLA on CD8+ TIL activated the PI3K-Akt pathway resulting in the phosphorylation of Akt; this response was blocked using an anti-BTLA antibody. PI3K-Akt activation is most likely mediated by a unique GRB2-binding domain that exists in BTLA that can recruit PI3K via GRB2. As HVEM is expressed by many melanoma tumors and antigen-presenting cells these results suggest that HVEM ligation of BTLA on CD8+BTLA+ effector T-cells in the tumor microenvironment may play a dual role by reducing over-stimulation through the TCR and driving a PI3K-Akt-induced cell survival pathway. This survival signaling pathway may facilitate the longer persistence of the CD8+BTLA+ TIL subset in vivo in our patients. Our study has uncovered a novel role for BTLA both as a biomarker in TIL therapy and as a rheostat in perhaps fine tuning CD8+ T-cell responses rather than simply as a negative signaling pathway.

Authors' details

¹MD Anderson, Houston, TX, USA. ²National Institutes of Health, Bethesda, MD, USA. ³Sidra Medical Research Hospital, Doha, Qatar.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-O1

Cite this article as: Haymaker et al.: Uncovering a novel function of BTLA on tumor-infiltrating CD8+ T cells. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):O1.

¹MD Anderson, Houston, TX, USA

Full list of author information is available at the end of the article